INDEPENDENT RESURVEY OF THE SUBDIVISIONAL LINES OF T. 28 N., R. 69 E.		
CHAINS		
54.40	Wash, 20 lks. wide, 1 ft. deep, drains E.	
79.65	The cor. of secs. 5, 6, 31 and 32, on the N. bdy. of the Tp., monumented with an iron post, $2\frac{1}{2}$ ins. diam., set, mkd. and witnessed as described in the field notes of the independent resurvey of the south boundary of T. 29 N., R. 69 E., executed under Group No. 463, in 1969.	
	Land, rolling. Soil, sandy clay and rocky. Timber, juniper and pinon. Undergrowth, sagebrush, greasewood and bunchgrass.	
	GENERAL DESCRIPTION	
	The land within Township 28 North, Range 69 East, ranges from about 5,100 to 8,000 ft. above sea level, and is rolling to mountainous. The soil is sandy clay, gravelly and rocky. The vegetation consists of sagebrush, greasewood and bunchgrass. There are scattered juniper and pinon on the higher elevations.	
,	The township is drained by numerous dry washes which wind northeasterly through the township.	
	No mineral formations of consequence were noted during the survey.	
	Principal users of the area are cattlemen. Access into the township is provided by desert trail roads.	
	54.40	CHAINS 54.40 Wash, 20 lks. wide, 1 ft. deep, drains E.  79.65 The cor. of secs. 5, 6, 31 and 32, on the N. bdy. of the Tp., monumented with an iron post, 2½ ins. diam., set, mkd. and witnessed as described in the field notes of the independent resurvey of the south boundary of T. 29 N., R. 69 E., executed under Group No. 463, in 1969.  Land, rolling. Soil, sandy clay and rocky. Timber, juniper and pinon. Undergrowth, sagebrush, greasewood and bunchgrass.  GENERAL DESCRIPTION  The land within Township 28 North, Range 69 East, ranges from about 5,100 to 8,000 ft. above sea level, and is rolling to mountainous. The soil is sandy clay, gravelly and rocky. The vegetation consists of sagebrush, greasewood and bunchgrass. There are scattered juniper and pinon on the higher elevations.  The township is drained by numerous dry washes which wind northeasterly through the township.  No mineral formations of consequence were noted during the survey.  Principal users of the area are cattlemen. Access into